

\*\*Volume Title\*\*

ASP Conference Series, Vol. \*\*Volume Number\*\*

\*\*Author\*\*

© \*\*Copyright Year\*\* Astronomical Society of the Pacific

## PHL 1811 Analogs: A Population of X-ray Weak Quasars

Jianfeng Wu,<sup>1</sup> W. N. Brandt,<sup>1</sup> P. B. Hall,<sup>2</sup> R. R. Gibson,<sup>3</sup> G. T. Richards,<sup>4</sup>  
D. P. Schneider,<sup>1</sup> O. Shemmer,<sup>5</sup> D. W. Just,<sup>6</sup> and S. J. Schmidt<sup>3</sup>

<sup>1</sup>*The Pennsylvania State University, University Park, PA 16802, USA*

<sup>2</sup>*York University, Toronto, ON M3J 1P3, Canada*

<sup>3</sup>*University of Washington, Seattle, WA 98915, USA*

<sup>4</sup>*Drexel University, Philadelphia, PA 19104, USA*

<sup>5</sup>*University of North Texas, Denton, TX 76203, USA*

<sup>6</sup>*Steward Observatory, Tucson, AZ 85721, USA*

**Abstract.** We report on a population of X-ray weak quasars with similar UV emission-line properties to those of the remarkable quasar PHL 1811. All radio-quiet PHL 1811 analogs are notably X-ray weak by a mean factor of  $\approx 13$ , with hints of heavy X-ray absorption. Correlations between the X-ray weakness and UV emission-line properties suggest that PHL 1811 analogs may have extreme wind-dominated broad emission-line regions (BELRs). We propose an AGN geometry that can potentially unify the PHL 1811 analogs and the general population of weak-line quasars.

Luminous X-ray emission is a nearly universal property of efficiently accreting supermassive black holes. However, there are a few examples of active galactic nuclei (AGNs) that emit X-rays much more weakly. An extreme case is the optically bright quasar PHL 1811 ( $z = 0.192$ ). It is consistently X-ray weak by a factor of  $\approx 30$ –100 compared to typical quasars with similar UV luminosity (Leighly et al. 2007a). This quasar also has unusual UV emission-line properties: weak and strongly blueshifted high-ionization lines (e.g., C iv), weak semi-forbidden lines (e.g., C iii) and forbidden lines (e.g., [O iii]), and strong UV Fe ii/iii emission (Leighly et al. 2007b).

To determine if there is a *population* of such X-ray weak quasars and to explore the connection between their UV emission-line properties and X-ray properties, we conducted a *Chandra* survey of a sample of Sloan Digital Sky Survey (York et al. 2000) quasars with similar UV emission-line properties to those of PHL 1811 (Wu et al. 2011a). Their relative X-ray brightness is quantified by the  $\Delta\alpha_{\text{ox}}$  parameter.<sup>1</sup> All radio-quiet PHL 1811 analogs, without exception, are remarkably X-ray weak by a mean factor of  $\approx 13$  (mean  $\Delta\alpha_{\text{ox}} = -0.42 \pm 0.05$ ; see Fig. 1a), which shows the connection between soft spectral energy distributions (X-ray weak and optical/UV strong) and PHL 1811-like UV emission lines. Their  $\Delta\alpha_{\text{ox}}$  values are correlated with the

---

<sup>1</sup> $\alpha_{\text{ox}}$  is defined to be the slope of an assumed power law connecting the rest-frame 2500 Å and 2 keV luminosities, i.e.,  $\alpha_{\text{ox}} = 0.384 \log(L_{2 \text{ keV}}/L_{2500 \text{ Å}})$ . This quantity is well known to be correlated with  $L_{2500 \text{ Å}}$  (e.g., Just et al. 2007; see Fig. 1a). We also define  $\Delta\alpha_{\text{ox}} = \alpha_{\text{ox}}(\text{observed}) - \alpha_{\text{ox}}(L_{2500 \text{ Å}})$ , which quantifies the observed X-ray luminosity relative to that expected from the  $\alpha_{\text{ox}}-L_{2500 \text{ Å}}$  relation.




Figure 1. (a):  $\alpha_{\text{ox}}-L_{2500 \text{ \AA}}$  diagram for radio-quiet PHL 1811 analogs (filled squares) and a comparison sample of typical radio-quiet quasars without broad absorption lines from the literature (open symbols). Downward arrows represent  $\alpha_{\text{ox}}$  upper limits. The solid line shows the  $\alpha_{\text{ox}}-L_{2500 \text{ \AA}}$  correlation in Just et al. (2007). (b): Schematic illustration of the WLQ/PHL 1811 analog unification hypothesis. Each panel is a side view of an accretion disk around a black hole. The X-ray continuum source is close to the black hole. Normally, the shielding gas only covers part of the BELR (top panel). When the shielding gas has a high BELR covering factor (bottom panel), it absorbs most of the X-ray photons, and a strong wind is generated in the BELR. When such a quasar is viewed through the shielding gas, an X-ray weak PHL 1811 analog is seen. When such a quasar is viewed away from the wind direction, an X-ray normal WLQ is seen.

C iv equivalent width and blueshift, showing our sample may represent quasars whose BELRs are dominated by AGN winds (see Richards et al. 2011). Our radio-quiet PHL 1811 analogs show a hard average X-ray spectrum ( $\Gamma = 1.1^{+0.45}_{-0.40}$ ) which may indicate heavy X-ray absorption. Observationally, the PHL 1811 analogs appear to be a subset ( $\approx 30\%$ ) of weak-line quasars (WLQs; e.g., Diamond-Stanic et al. 2009; Shemmer et al. 2009). We propose an AGN geometry that can potentially unify the PHL 1811 analogs and the general WLQ population via orientation effects (see Fig. 1b). This model assumes a subset of quasars in which the highly ionized “shielding gas” (e.g., Murray et al. 1995) covers most of the BELR and blocks the X-ray photons from reaching the BELR, resulting in weak high-ionization lines and strong wind acceleration. Our recent work on low-redshift WLQs (Wu et al. 2011b) provides support to this unification model.

## References

- Diamond-Stanic, A. M., Fan, X., Brandt, W. N., et al. 2009, *ApJ*, 699, 782  
 Gibson, R. R., Brandt, W. N., & Schneider, D. P. 2008, *ApJ*, 685, 773  
 Just, D. W., Brandt, W. N., Shemmer, O., et al. 2007, *ApJ*, 665, 1004  
 Leighly, K. M., Halpern, J. P., Jenkins, E. B., et al. 2007a, *ApJ*, 663, 103  
 Leighly, K. M., Halpern, J. P., Jenkins, E. B., & Casebeer, D. 2007b, *ApJS*, 173, 1  
 Murray, N., Chiang, J., Grossman, S. A., & Voit, G. M. 1995, *ApJ*, 451, 498  
 Richards, G. T., Kruczek, N. E., Gallagher, S. C., et al. 2011, *AJ*, 141, 167  
 Shemmer, O., Brandt, W. N., Schneider, D. P., et al. 2006, *ApJ*, 644, 86  
 Shemmer, O., Brandt, W. N., Anderson, S. F., et al. 2009, *ApJ*, 696, 580  
 Wu, J., Brandt, W. N., Hall, P. B., et al. 2011a, *ApJ*, 736, 28  
 Wu, J., Brandt, W. N., Anderson, S. F., et al. 2011b, *ApJ* in press (arXiv:1112.2228 [astro-ph])  
 York, D. G., Adelman, J., Anderson, J. E., Jr., et al. 2000, *AJ*, 120, 1579